

--	--	--	--	--

--	--	--	--

9701/02

May/June 2007

1 hour 15 minutes

Additional Materials: Data Booklet

READ THESE INSTRUCTIONS FIRST

DO **NOT** WRITE IN ANY BARCODES.

At the end of the examination, fasten all your work securely together.

DO NOT WRITE IN THE GREY AREAS BETWEEN THE PAGES.

For Examiner's Use	
1	
2	
3	
4	
Total	

This document consists of **9** printed pages and **3** blank pages.

Answer **all** the questions in the spaces provided.

For
Examiner's
Use

- 1 Ethene, C_2H_4 , and hydrazine, N_2H_4 , are hydrides of elements which are adjacent in the Periodic Table. Data about ethene and hydrazine are given in the table below.

	C_2H_4	N_2H_4
melting point/ $^{\circ}\text{C}$	-169	+2
boiling point/ $^{\circ}\text{C}$	-104	+114
solubility in water	insoluble	high
solubility in ethanol	high	high

- (a) Ethene and hydrazine have a similar arrangement of atoms but differently shaped molecules.

- (i) What is the H-C-H bond angle in ethene?

.....

- (ii) Draw a 'dot-and-cross' diagram for hydrazine.

- (iii) What is the H-N-H bond angle in hydrazine?

.....

[4]

- (b) The melting and boiling points of hydrazine are much higher than those of ethene. Suggest reasons for these differences in terms of the intermolecular forces **each** compound possesses.

.....

.....

.....

.....

.....

[3]

- (c) Explain, with the aid of a diagram showing lone pairs of electrons and dipoles, why hydrazine is very soluble in ethanol.

For
Examiner's
Use

[3]

Ethene and hydrazine each react with HCl .

- (d) When ethene is reacted with HCl , $\text{C}_2\text{H}_5\text{Cl}$ is the only product.

- (i) Using structural formulae, give an equation for the reaction between ethene and HCl .

- (ii) What type of reaction occurs between HCl and ethene?

.....

- (iii) Explain why there is no further reaction between $\text{C}_2\text{H}_5\text{Cl}$ and HCl .

.....

[3]

- (e) When aqueous hydrazine is reacted with HCl , a solid compound of formula $\text{N}_2\text{H}_5\text{Cl}$ may be isolated. When an excess of HCl is used, a second solid, $\text{N}_2\text{H}_6\text{Cl}_2$, is formed.

- (i) Suggest what type of reaction occurs between hydrazine and HCl .

.....

- (ii) What feature of the hydrazine molecule enables this reaction to occur?

.....

- (iii) Suggest why one molecule of hydrazine is able to react with one or two molecules of HCl .


.....

.....

[3]

[Total: 16]

- 2 Alcohols and esters are important organic compounds which are widely used as solvents. Esters such as ethyl ethanoate can be formed by reacting carboxylic acids with alcohols.



This reaction is an example of a dynamic equilibrium.

- (a) Explain what is meant by the term *dynamic equilibrium*.

.....
[1]

- (b) Write the expression for the equilibrium constant for this reaction, K_c .

[1]

- (c) For this equilibrium, the value of K_c is 4.0 at 298 K.

A mixture containing 0.5 mol of ethanoic acid, 0.5 mol ethanol, 0.1 mol ethyl ethanoate and 0.1 mol water was set up and allowed to come to equilibrium at 298 K. The final volume of solution was $V \text{ dm}^3$.

Calculate the amount, in moles, of each substance present at equilibrium.

[4]

Alcohols may be classified into primary, secondary and tertiary. Some reactions are common to all three types of alcohol. In other cases, the same reagent gives different products depending on the nature of the alcohol.

For
Examiner's
Use

- (d) In the empty squares below give the structural formula of the organic compound formed in each of the reactions indicated.

If no reaction occurs, write 'no reaction' in the space.

alcohol reagent(s) and conditions	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$	$\text{CH}_3\text{CH}_2\text{CH}(\text{OH})\text{CH}_3$	$(\text{CH}_3)_3\text{COH}$
red phosphorus and iodine heat under reflux			
concentrated H_2SO_4 heat			
$\text{Cr}_2\text{O}_7^{2-}/\text{H}^+$ heat under reflux			

[5]

[Total: 11]

- 3** This question is about the elements in Group II of the Periodic Table, magnesium to barium.

For
Examiner's
Use

- (a)** Complete the table below to show the electronic configuration of calcium atoms and of strontium ions, Sr^{2+} .

	1s	2s	2p	3s	3p	3d	4s	4p	4d
Ca	2	2	6						
Sr^{2+}	2	2	6						

[2]

- (b)** Explain the following observations.

- (i)** The atomic radii of Group II elements increase down the Group.

.....

- (ii)** The strontium ion is smaller than the strontium atom.

.....

- (iii)** The first ionisation energies of the elements of Group II decrease with increasing proton number.

.....

[4]

- (c) Samples of magnesium and calcium are placed separately in cold water and left for some time. In **each case**, describe what you would see and write a balanced equation for each reaction.

(i) magnesium

observation

.....

equation

(ii) calcium

observation

.....

equation

[6]

- (d) Strontium nitrate, $\text{Sr}(\text{NO}_3)_2$ undergoes thermal decomposition.

(i) State one observation you would make during this reaction.

.....

.....

(ii) Write a balanced equation for this reaction.

.....

[4]

[Total: 16]

- 4 Commercial paint and varnish removers contain a mixture of dichloromethane, CH_2Cl_2 , and methanol, CH_3OH .

(a) What would be observed when the following reactions are carried out?
In each case, give the name or formula of the reaction product which is responsible for the observation you have made.

- (i) CH_2Cl_2 is reacted with $\text{NaOH}(\text{aq})$ and $\text{AgNO}_3(\text{aq})$ and the mixture left to stand.

observation

product responsible

- (ii) CH_3OH is mixed with PCl_5 .

observation

product responsible

- (iii) CH_3OH is reacted with sodium.

observation

product responsible

[6]

- (b) When CH_2Cl_2 is heated under reflux with an excess of $\text{NaOH}(\text{aq})$, a compound **W** is formed.

W has the following composition by mass: C, 40.0%; H, 6.7%; O, 53.3%.

Use this information and the *Data Booklet* to show that the empirical formula of **W** is CH_2O .

[2]

- (c) Compounds with the empirical formula CH_2O can have the molecular formula $\text{C}_2\text{H}_4\text{O}_2$.

Two possible structural formulae for compounds with molecular formula $\text{C}_2\text{H}_4\text{O}_2$ are HCO_2CH_3 and $\text{H}_2\text{C}=\text{C}(\text{OH})_2$.

In the boxes below, draw displayed formulae for **three further** structural isomers with the molecular formula $\text{C}_2\text{H}_4\text{O}_2$.

Do **not** attempt to draw any structures containing rings or O–O bonds.

X	Y	Z
----------	----------	----------

[3]

- (d) Identify which of your compounds, **X**, **Y**, or **Z**, will react with the following reagents.

In **each** case, state what you would observe.

- (i) solid NaHCO_3

compound

observation

- (ii) Tollens' reagent

compound

observation

[4]

- (e) **One** of the three compounds, **X**, **Y**, or **Z**, shows stereoisomerism.

Draw displayed, labelled structures of the stereoisomers of this compound.

[2]

[Total: 17]

BLANK PAGE

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.